

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
15 December 2005 (15.12.2005)

PCT

(10) International Publication Number
WO 2005/118561 A1

(51) International Patent Classification⁷: **C07D 263/58**,
A61K 31/423, A61P 25/16

(21) International Application Number:
PCT/EP2005/005596

(22) International Filing Date: 24 May 2005 (24.05.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
04012573.4 27 May 2004 (27.05.2004) EP

(71) Applicant (*for all designated States except US*): **UCB, S.A.**
[BE/BE]; Allée de la recherche 60, B-1070 Bruxelles (BE).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **STARCK,**
Jean-Philippe [FR/FR]; Rue des Tilleuls 12, F-67270
Gougenheim (FR). **KENDA, Benoît** [BE/BE]; Rue du
Hazoir, 41A, B-5080 Emnes (BE).

(74) Agent: **LECHIEN, Monique**; UCB, S.A., Allée de la
Recherche 60, B-1070 Bruxelles (BE).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:
— *with international search report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: BENZOXAZOLONE DERIVATIVES, PROCESSES FOR PREPARING THEM AND THEIR USES

(57) Abstract: The present invention relates to benzoxazolone derivatives, processes for preparing them, pharmaceutical compositions containing them and their use as pharmaceuticals, in the treatment of movement disorders, in particular in Parkinson's disease.



BENZOXAZOLONE DERIVATIVES, PROCESSES FOR PREPARING THEM AND THEIR USES

The present invention concerns benzoxazolone derivatives, processes for
5 preparing them, pharmaceutical compositions containing them and their use as
pharmaceuticals.

Movement and other disorders due to dysfunction of the basal ganglia and
related brain structures are of major socio-economic importance. Such disorders can
occur as a consequence of inherited or acquired disease, idiopathic neurodegeneration
10 or they may be iatrogenic. The spectrum of disorders is very diverse, ranging from
those associated with poverty of movement (akinesia, hypokinesia, bradykinesia, e.g.
in parkinsonian symptomatology), hypertonia (e.g. Parkinson's disease, spasticity) to
the involuntary movement disorders (hyperkinesias or dyskinesia, e.g. Huntington's
disease, dyskinesia induced by L-3,4-dihydroxyphenylalanine (L-DOPA or levodopa),
15 tardive dyskinesia, progressive supranuclear palsy, multiple system atrophy,
corticobasal degeneration, Wilson's disease, progressive pallidal atrophy, the
dystonias, metabolic neurotransmitter diseases, tics, tremor, Tourette's syndrome,
Sydenham's chorea, restless legs syndrome (RLS), chorea and choreathetosis,
paroxysmal dyskinesias, myoclonic disorders, Rett syndrome).

20 Parkinson's disease and related conditions are amongst of the most prevalent
diseases associated with poverty of movement. Parkinsonian symptoms are
characterized by slowness of movement (bradykinesia), rigidity and/or tremor.
Parkinsonian symptoms are seen in a variety of conditions, most commonly in
idiopathic parkinsonism (i.e. Parkinson's Disease) but also following treatment of
25 schizophrenia (i.e. neuroleptic-induced parkinsonism), exposure to toxins/drugs and
head injury.

It is widely appreciated that the primary pathology underlying Parkinson's
disease is degeneration, in the brain, of the dopaminergic projections from the
substantia nigra to the striatum. This has led to the widespread use of dopamine-
30 replacing agents (e.g. L-3,4-dihydroxyphenylalanine (L-DOPA) and dopamine agonists)
as symptomatic treatments for Parkinson's disease .

L-DOPA therapy currently offers the best symptomatic treatment of
Parkinson's disease and a variety of other movement disorders and most patients will
require it during the course of their disease. However, patients will develop L-DOPA-
35 associated motor complications. Problems can include motor fluctuations (e.g. delayed
"on" response and dose failure, end-of-dose wearing-off, unpredictable "on-off"
response, freezing episodes) and the appearance of a range of side-effects which
manifest as abnormal involuntary movements, such as dyskinesia (e.g. peak-dose

dyskinesia, "off" dystonia, diphasic dyskinesia). Dyskinesias are usually dystonic or choreiform in nature.

The phenomenon of "end-of-dose wearing-off" generally occurs early in the course of the disease. This is the most common and usually the first type of motor fluctuation that develops. As the name implies, the patient develops a loss of response to a dose of medication before taking the next dose. This occurs more often with levodopa than with the dopamine agonists because the agonists have a significantly longer half-life. Over time, fluctuation from mobility to immobility occurs more frequently, becoming more abrupt and disabling. The response to treatment can become unpredictable, many doses of levodopa having a delayed effect or even no effect at all.

Although many attempts have been made to develop agents that will prevent the development an/or the expression of dyskinesias, just a few were made to find a therapeutic tool able to manage motor fluctuations. Until now, only two classes of drug compounds, the catechol-O-methyltransferase (COMT) and monoamine oxydase type B (MAO_B) inhibitors, were developed to lengthen the beneficial therapeutic effect of L-DOPA but some of these compounds show adverse effects.

There is therefore, a need for new add-on therapies to L-DOPA which can enhance its efficacy and/or reduce its adverse effects.

Acta Poloniae Pharm. 1989, 46(2), 114-118, discloses 2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide, 5-chloro-2-oxo-3(2H)-benzoxazoleacetamide and 6-chloro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide as synthesis intermediates in the synthesis of N-carbamoylmethyl derivatives.

J. Prakt. Chem. 1966, 33(3-4), 130-138, discloses 2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide as a synthesis intermediate in the synthesis of (benzoxazol-3-yl)-carboxylic acids.

Godishnik na Sofiiskiya Universitet Sv. Kliment Okhridski, Khimicheski Fakultet (1979), Volume Date 1975-1976, 70, Pt. 2, 35-9, discloses N-methyl-2-oxo-3(2H)-benzoxazoleacetamide.

Khimiya Geterostsiklicheskikh Soedinenii (1985), (9), 1185-8, discloses: 6-bromo-5-chloro-N-methyl-2-oxo-3(2H)-benzoxazoleacetamide; 6-bromo-N-methyl-2-oxo-3(2H)-benzoxazoleacetamide; 6-chloro-N-methyl-2-oxo-3(2H)-benzoxazoleacetamide; 5-chloro-N-methyl-2-oxo-3(2H)-benzoxazoleacetamide and N-methyl-2-oxo-3(2H)-benzoxazoleacetamide.

Uzbekskii Khimicheskii Zhurnal (1988), (1), 70 discloses : N-methyl-2-oxo-3(2H)-benzoxazoleacetamide and N-(1,1-dimethylethyl)-2-oxo-3(2H)-benzoxazoleacetamide.

Zhurnal Vsesoyuznogo Khimicheskogo Obshchestva im. D.I. Mendeleeva (1987), 32(3), 342-4, discloses : 6-bromo-N,N-diethyl-2-oxo-3(2H)-benzoxazoleacetamide; 6-chloro-N,N-diethyl-2-oxo-3(2H)-benzoxazoleacetamide; N,N-diethyl-2-oxo-3(2H)-benzoxazoleacetamide; 6-bromo-N,N-dibutyl-2-oxo-3(2H)-benzoxazoleacetamide; N,N-dibutyl-6-chloro-2-oxo-3(2H)-benzoxazoleacetamide and N,N-dibutyl-2-oxo-3(2H)-benzoxazoleacetamide.

Database Crossfire Beilstein Institute for Organic Chemistry, XP002300055 (BRN:4485565) describes 6-bromo-5-chloro-3-(methylcarbamoylmethyl)-2-benzoxazolone.

The following compounds are disclosed in chemical libraries: 5-bromo-2-oxo-3(2H)-benzoxazoleacetamide; 6-bromo-2-oxo-3(2H)-benzoxazoleacetamide; 2-oxo-N-propyl-3(2H)-benzoxazoleacetamide; N,N-bis(2-methylpropyl)-6-chloro-2-oxo-3(2H)-benzoxazoleacetamide; 6-chloro-2-oxo-N-propyl-3(2H)-benzoxazoleacetamide; N-(1-methylethyl)-2-oxo-3(2H)-benzoxazoleacetamide; 6-chloro-N-(1-methylethyl)-2-oxo-3(2H)-benzoxazoleacetamide; 6-chloro-N-(1,1-dimethylethyl)-2-oxo-3(2H)-benzoxazoleacetamide; 5-chloro-2-oxo-1,3-benzoxazol-3(2H)-yl-N-(1,1-dimethylethyl)acetamide; 2-oxo-1,3-benzoxazol-3(2H)-yl-N-(1-methylpropyl)acetamide; 6-chloro-N,N-dimethyl-2-oxo-3(2H)-benzoxazoleacetamide and N,N-dimethyl-2-oxo-3(2H)-benzoxazoleacetamide.

It is known from the international patent application WO 00/29397 that piperazine and piperidine derivatives have anti-anxiety and anti-depression activities.

It is known from the international patent application WO 94/29272 that 1-substituted isatin and oxindole derivatives are of value in the treatment of the cognitive dysfunctions.


It is known from the international patent application WO 03/076420 that cyclic amide derivatives are of value in the treatment of neurodegenerative diseases.

It is known from the European patent application EP 0 673 933 that aminoalkyl benzoxazolinone and benzothiazolinone derivatives have anti-anxiety activity and psychotropic activity.

It has now surprisingly been found that certain benzoxazolone derivatives demonstrate therapeutic properties which render them useful in a variety of pharmaceutical indications, and particularly for the symptomatic and/or prophylactic treatment of movement disorders and/or motor fluctuations, in particular in Parkinson's disease.

In one aspect the invention therefore provides a compound having formula I or a pharmaceutically acceptable salt thereof,

4



wherein

R¹ is hydrogen or C₁₋₄-alkyl unsubstituted or substituted by hydroxy;

R² is hydrogen or C₁₋₄-alkyl unsubstituted or substituted by hydroxy;

5 R³ is hydrogen or unsubstituted C₁₋₄-alkyl;

R^{3a} is hydrogen or unsubstituted C₁₋₄-alkyl;

R⁴ is hydrogen, halogen, or C₁₋₄-alkoxy unsubstituted or substituted by a phenyl group;

R⁵ is hydrogen, halogen, or C₁₋₄-alkoxy unsubstituted or substituted by a phenyl group;

10 R⁶ is hydrogen, halogen, or C₁₋₄-alkoxy unsubstituted or substituted by a phenyl group;

R⁷ is hydrogen, halogen, or C₁₋₄-alkoxy unsubstituted or substituted by a phenyl group;

with the proviso that if R¹, R², R⁴, R⁵ and R⁷ are hydrogen, then R⁶ is not hydrogen, bromo or chloro; and if R¹, R², R⁴, R⁶ and R⁷ are hydrogen, then R⁵ is not bromo or chloro; and if R¹, R², R³, R⁴ and R⁷ are hydrogen, R⁶ is bromo, and R⁵ is chloro, then R^{3a} is not methyl.

15 In another aspect the invention provides 2-(6-bromo-2-oxo-1,3-benzoxazol-3(2H)-yl)-N,N-dimethylacetamide.

20 The term "halogen", as used herein, includes chloro, bromo, fluoro, iodo. Preferred halogens are chloro, bromo, and fluoro.

The term "hydroxy", as used herein, represents a group of formula -OH.

The term "C₁₋₄-alkoxy", as used herein, represents a group of formula -OR^a

25 wherein R^a is a C₁₋₄-alkyl group, optionally substituted by 1 to 3 substituents selected from halogen or a phenyl group. The preferred C₁₋₄-alkoxy group is methoxy.

The term "C₁₋₄-alkyl", as used herein, is defined as including saturated, monovalent hydrocarbon radicals having straight or branched moieties and containing 1-4 carbon atoms. Any alkyl moiety may optionally be substituted by 1 to 5

30 substituents selected independently from hydroxy, halogen or a phenyl group. The preferred alkyl group is methyl.

The term "phenyl group", as used herein, represents an organic radical derived from an aromatic hydrocarbon consisting of a ring and containing 6 carbon atoms by

removal of one hydrogen, and optionally substituted by 1 to 3 substituents selected from halogen, C₁₋₄-alkyl, C₁₋₄-alkoxy or cyano.

The term "cyano" as used herein, represents a group of the formula -CN.

Generally, R¹ is hydrogen or C₁₋₄-alkyl unsubstituted or substituted by
5 hydroxy. Preferred R¹ is hydrogen.

Generally, R² is hydrogen or C₁₋₄-alkyl unsubstituted or substituted by hydroxy. Preferred R² is hydrogen.

Generally, R³ is hydrogen or unsubstituted C₁₋₄-alkyl. Preferred R³ is hydrogen or methyl. More preferred R³ is hydrogen.

10 Generally, R^{3a} is hydrogen or unsubstituted C₁₋₄-alkyl. Preferred R^{3a} is hydrogen or methyl. More preferred R^{3a} is hydrogen.

Generally, R⁴ is hydrogen, halogen, or C₁₋₄-alkoxy unsubstituted or substituted by a phenyl group. Usually, R⁴ is hydrogen and halogen. Preferred R⁴ is hydrogen and chloro. More preferred R⁴ is hydrogen.

15 Generally, R⁵ is hydrogen, halogen, or C₁₋₄-alkoxy unsubstituted or substituted by a phenyl group. Usually, R⁵ is hydrogen and halogen. Preferred R⁵ is hydrogen and fluoro. More preferred R⁵ is hydrogen.

Generally, R⁶ is hydrogen, halogen, or C₁₋₄-alkoxy unsubstituted or substituted by a phenyl group. Usually, R⁶ is hydrogen, halogen or unsubstituted C₁₋₄-alkoxy. Preferred is hydrogen, fluoro, bromo, chloro and methoxy. More preferred
20 R⁶ is hydrogen, fluoro, chloro and methoxy.

Generally, R⁷ is hydrogen, halogen, or C₁₋₄-alkoxy unsubstituted or substituted by a phenyl group. Usually, R⁷ is hydrogen, halogen or unsubstituted C₁₋₄-alkoxy. Preferred R⁷ is hydrogen, fluoro, bromo, chloro and methoxy. More preferred
25 R⁷ is hydrogen, fluoro, bromo and chloro.

Combinations of one or more of these preferred compound groups are especially preferred.

In another aspect the invention therefore provides a compound having formula I or a pharmaceutically acceptable salt thereof, with the proviso that if R¹, R², R⁴, R⁵
30 and R⁷ are hydrogen, then R⁶ is not hydrogen, bromo or chloro; and if R¹, R², R⁴, R⁶ and R⁷ are hydrogen, then R⁵ is not bromo or chloro; and if R¹, R², R⁴ and R⁷ are hydrogen, and R⁶ is bromo, then R⁵ is not chloro.

In another aspect the invention therefore provides a compound having formula I or a pharmaceutically acceptable salt thereof, with the proviso that if R¹, R², R⁴ and
35 R⁷ are hydrogen, then R⁵ and/or R⁶ are not hydrogen, bromo or chloro.

In one embodiment of the invention, the compounds of formula I, or pharmaceutically acceptable salts thereof, are those wherein

R¹ is hydrogen;

R² is hydrogen;

R³ is selected from hydrogen or unsubstituted C₁₋₄-alkyl;

R^{3a} is selected from hydrogen or unsubstituted C₁₋₄-alkyl;

5 R⁴ is selected from hydrogen or halogen;

R⁵ is selected from hydrogen or halogen;

R⁶ is selected from hydrogen, halogen or unsubstituted C₁₋₄-alkoxy;

R⁷ is selected from hydrogen, halogen or unsubstituted C₁₋₄-alkoxy;

10 with the proviso that if R⁴, R⁵ and R⁷ are hydrogen, then R⁶ is not hydrogen, bromo or chloro; and if R⁴, R⁶ and R⁷ are hydrogen, then R⁵ is not bromo or chloro; and if R³, R⁴ and R⁷ are hydrogen, R⁶ is bromo, and R⁵ is chloro, then R^{3a} is not methyl.

In a preferred embodiment of the invention, the compounds of formula I, or pharmaceutically acceptable salts thereof, are those wherein

15 R¹ is hydrogen;

R² is hydrogen;

R³ is selected from hydrogen or methyl;

R^{3a} is selected from hydrogen or methyl;

R⁴ is hydrogen or chloro;

20 R⁵ is hydrogen or fluoro;

R⁶ is selected from hydrogen, fluoro, chloro, bromo or methoxy;

R⁷ is selected from hydrogen, fluoro, chloro, bromo or methoxy;

25 with the proviso that if R⁴, R⁵ and R⁷ are hydrogen, then R⁶ is not hydrogen, bromo or chloro; and if R⁴, R⁶ and R⁷ are hydrogen, then R⁵ is not bromo or chloro; and if R³, R⁴ and R⁷ are hydrogen, R⁶ is bromo, and R⁵ is chloro, then R^{3a} is not methyl.

In a more preferred embodiment of the invention, the compounds of formula I, or pharmaceutically acceptable salts thereof, are those wherein

R¹ is hydrogen;

30 R² is hydrogen;

R³ is selected from hydrogen or methyl;

R^{3a} is selected from hydrogen or methyl;

R⁴ is hydrogen;

R⁵ is hydrogen;

35 R⁶ is selected from hydrogen, fluoro, chloro, or methoxy;

R⁷ is selected from hydrogen, fluoro, chloro, or bromo;

with the proviso that if R⁷ is hydrogen, then R⁶ is not hydrogen or chloro.

Preferred individual compounds of the invention are:

2-(6-methoxy-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
2-(6-fluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
2-(7-fluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
5 2-(5,6-difluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
2-(6-bromo-7-fluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
2-(7-chloro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
2-(6-bromo-7-chloro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
2-(6,7-dichloro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
10 2-(6,7-difluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
2-(6-chloro-7-fluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
2-(7-bromo-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
2-(6,7-dibromo-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
2-(7-bromo-6-chloro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
15 2-(7-methoxy-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
2-(4-chloro-6,7-difluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
2-(6-bromo-2-oxo-1,3-benzoxazol-3(2H)-yl)-N,N-dimethylacetamide.

More preferred compounds of the invention:

2-(6-methoxy-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
20 2-(6-fluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
2-(7-fluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
2-(7-chloro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
2-(7-bromo-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide; and
2-(6,7-difluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide.

Most preferred compounds of the invention:

2-(6-methoxy-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
2-(6-fluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
2-(7-fluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
2-(7-bromo-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide; and
30 2-(6,7-difluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide.

The "pharmaceutically acceptable salts" according to the invention include therapeutically active, non-toxic base salt forms which the compounds of formula I are able to form.

For example the compounds of formula I containing acidic protons may be
35 converted into their therapeutically active, non-toxic base addition salt forms, e.g. metal or amine salts, by treatment with appropriate organic and inorganic bases. Appropriate base salt forms include, for example but not limited to, ammonium salts,

alkali and alkaline earth metal salts, e.g. lithium, sodium, potassium, magnesium, calcium salts and the like, salts with organic bases, e.g. N-methyl-D-glucamine, hydrabamine salts, and salts with amino acids such as, for example, arginine, lysine and the like.

5 Conversely said salt forms can be converted into the free forms by treatment with an appropriate acid.

Compounds of the formula I and their salts can be in the form of solvates, which are included within the scope of the present invention. Such solvates include for example hydrates, alcoholates and the like.

10 Some of the compounds of formula I and some of their intermediates have at least one stereogenic center in their structure. This stereogenic center may be present in a R or a S configuration, said R and S notation is used in correspondence with the rules described in Pure Appl. Chem., 45 (1976) 11-30.

The invention also relates to all stereoisomeric forms such as enantiomeric and 15 diastereoisomeric forms of the compounds of formula I or mixtures thereof (including all possible mixtures of stereoisomers).


Some of the compounds of formula I may also exist in tautomeric forms. Such forms although not explicitly indicated in the above formula are intended to be included within the scope of the present invention.

20 With respect to the present invention reference to a compound or compounds is intended to encompass that compound in each of its possible isomeric forms and mixtures thereof unless the particular isomeric form is referred to specifically.

Compounds according to the present invention may exist in different 25 polymorphic forms. Although not explicitly indicated in the above formula, such forms are intended to be included within the scope of the present invention.

The compounds of formula I according to the invention can be prepared analogously to conventional methods as understood by the person skilled in the art of synthetic organic chemistry.


30 According to one embodiment, some compounds having the general formula I may be prepared by alkylation of a compound of formula II with a compound of formula III according to the equation:



wherein Hal is an halogen atom, preferably bromo or chloro.


This reaction may be carried out in the presence of a strong base, preferably sodium hydride, at a temperature between 0 and 40 °C, in an inert solvent, for example DMF, under an inert atmosphere, or as described in the British patent GB 1,309,692.

Compounds of formula II may be prepared by reaction of a compound of formula IV with 1,1'-carbonyldiimidazole of formula V according to the equation



This reaction may be carried out in an inert solvent at a temperature comprised between 0 and 100 °C.

Compounds of formula IV may be prepared by reduction of a compound of formula VI according to procedures known to the person skilled in the art.



Compounds of formula VI are available either commercially or using procedures known to the person skilled in the art.

According to another embodiment, some compounds having the general formula I may be prepared by halogenation of the corresponding compound of formula I wherein R⁶ is a hydrogen atom with a N-halosuccinimide or any suitable halogenation agent according to the procedure described in: Castanet A.-S., Colobert F., Broutin P.-E., Tetrahedron Lett. (2002), 43, 5047-5048.


According to another embodiment, some compounds having the general formula I may be prepared by halogenation of the corresponding compound of formula I wherein R⁴ is a hydrogen atom and R⁶ is different from hydrogen with a N-halosuccinimide or any suitable halogenation agent according to the procedure

described in: Castanet A.-S., Colobert F., Broutin P.-E., Tetrahedron Lett. (2002), 43, 5047-5048.

In one embodiment, the present invention concerns also the synthesis intermediates selected from the group consisting of 5,6-difluoro-1,3-benzoxazol-2(3H)-one and 7-bromo-1,3-benzoxazol-2(3H)-one.

It has now been found that compounds of formula Ia, their pharmaceutically acceptable salts, or stereoisomeric forms thereof

It has now been found that compounds of formula Ia, their pharmaceutically acceptable salts, or stereoisomeric forms thereof



wherein

R^1 is hydrogen or C_{1-4} -alkyl unsubstituted or substituted by hydroxy;

R^2 is hydrogen or C_{1-4} -alkyl unsubstituted or substituted by hydroxy;

R^3 is hydrogen or unsubstituted C_{1-4} -alkyl;

R^{3a} is hydrogen or unsubstituted C_{1-4} -alkyl;

R^4 is hydrogen, halogen, or C_{1-4} -alkoxy unsubstituted or substituted by a phenyl group;

R^5 is hydrogen, halogen, or C_{1-4} -alkoxy unsubstituted or substituted by a phenyl group;


R^6 is hydrogen, halogen, or C_{1-4} -alkoxy unsubstituted or substituted by a phenyl group;

R^7 is hydrogen, halogen, or C_{1-4} -alkoxy unsubstituted or substituted by a phenyl group;

are useful in a variety of pharmaceutical indications.

For example, the compounds of formula Ia according to the invention are useful for the symptomatic and/or prophylactic treatment of movement disorders and/or motor fluctuations, in particular in Parkinson's disease.

In another aspect the invention therefore provides the therapeutical use of compounds of formula Ia, or pharmaceutically acceptable salts thereof,



wherein

R^1 is hydrogen or C_{1-4} -alkyl unsubstituted or substituted by hydroxy;

R^2 is hydrogen or C_{1-4} -alkyl unsubstituted or substituted by hydroxy;

5 R^3 is hydrogen or unsubstituted C_{1-4} -alkyl;

R^{3a} is hydrogen or unsubstituted C_{1-4} -alkyl;


R^4 is hydrogen, halogen, or C_{1-4} -alkoxy unsubstituted or substituted by a phenyl group;

10 R^5 is hydrogen, halogen, or C_{1-4} -alkoxy unsubstituted or substituted by a phenyl group;

R^6 is hydrogen, halogen, or C_{1-4} -alkoxy unsubstituted or substituted by a phenyl group;

R^7 is hydrogen, halogen, or C_{1-4} -alkoxy unsubstituted or substituted by a phenyl group.

15 In a particular embodiment, the invention provides the therapeutical use of compounds of formula I, or pharmaceutically acceptable salts thereof,



wherein

20 R^1 is hydrogen or C_{1-4} -alkyl unsubstituted or substituted by hydroxy;

R^2 is hydrogen or C_{1-4} -alkyl unsubstituted or substituted by hydroxy;

R^3 is hydrogen or unsubstituted C_{1-4} -alkyl;

R^{3a} is hydrogen or unsubstituted C_{1-4} -alkyl;

25 R^4 is hydrogen, halogen, or C_{1-4} -alkoxy unsubstituted or substituted by a phenyl group;

R^5 is hydrogen, halogen, or C_{1-4} -alkoxy unsubstituted or substituted by a phenyl group;

R^6 is hydrogen, halogen, or C_{1-4} -alkoxy unsubstituted or substituted by a phenyl group;

R⁷ is hydrogen, halogen, or C₁₋₄-alkoxy unsubstituted or substituted by a phenyl group;

with the proviso that if R¹, R², R⁴, R⁵ and R⁷ are hydrogen, then R⁶ is not hydrogen, bromo or chloro; and if R¹, R², R⁴, R⁶ and R⁷ are hydrogen, then R⁵ is not bromo or chloro; and if R¹, R², R³, R⁴ and R⁷ are hydrogen, R⁶ is bromo, and R⁵ is chloro, then R^{3a} is not methyl.

The invention also provides the therapeutical use of the compound 2-(6-bromo-2-oxo-1,3-benzoxazol-3(2H)-yl)-N,N-dimethylacetamide, or of its pharmaceutically acceptable salts.

In a preferred embodiment, the invention provides the therapeutical use of compounds of formula Ia, or pharmaceutically acceptable salts thereof, wherein

R¹ is hydrogen;

R² is hydrogen;

R³ is selected from hydrogen or unsubstituted C₁₋₄-alkyl;

R^{3a} is selected from hydrogen or unsubstituted C₁₋₄-alkyl;

R⁴ is hydrogen;

R⁵ is selected from hydrogen, bromo or chloro;

R⁶ is selected from hydrogen, bromo or chloro and

R⁷ is hydrogen.


In a preferred embodiment, the invention provides the therapeutical use of a compound of formula Ia selected from 2-(6-bromo-2-oxo-1,3-benzoxazol-3(2H)-yl)-N,N-dimethylacetamide; 2-(2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide; 2-(6-bromo-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide; 2-(6-chloro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide; 2-(6-chloro-2-oxo-1,3-benzoxazol-3(2H)-yl)-N,N-dimethylacetamide; and N,N-dimethyl-2-(2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide.

In a more preferred embodiment, the invention provides the therapeutical use of a compound of formula Ia selected from 2-(6-bromo-2-oxo-1,3-benzoxazol-3(2H)-yl)-N,N-dimethylacetamide; 2-(2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide; and 2-(6-chloro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide.

In a most preferred embodiment, the invention provides the therapeutical use of a compound of formula Ia selected from 2-(2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide and 2-(6-chloro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide.

In another embodiment, the invention provides a pharmaceutical composition comprising a compound of formula Ia, its pharmaceutically acceptable salts, or stereoisomeric form thereof

13



wherein

R¹ is hydrogen or C₁₋₄-alkyl unsubstituted or substituted by hydroxy;

R² is hydrogen or C₁₋₄-alkyl unsubstituted or substituted by hydroxy;

5 R³ is hydrogen or unsubstituted C₁₋₄-alkyl;

R^{3a} is hydrogen or unsubstituted C₁₋₄-alkyl;

R⁴ is hydrogen, halogen, or C₁₋₄-alkoxy unsubstituted or substituted by a phenyl group;

R⁵ is hydrogen, halogen, or C₁₋₄-alkoxy unsubstituted or substituted by a

10 phenyl group;

R⁶ is hydrogen, halogen, or C₁₋₄-alkoxy unsubstituted or substituted by a phenyl group;

R⁷ is hydrogen, halogen, or C₁₋₄-alkoxy unsubstituted or substituted by a phenyl group.

15 In another aspect the invention provides the use of compounds of formula Ia, or pharmaceutically acceptable salts thereof, for the symptomatic and/or prophylactic treatment of motor fluctuations and/or dyskinesia in Parkinson's patients before or during exposure to dopamine replacement therapy.

The compounds of formula Ia according to the invention may also be used for
20 the treatment and the prevention of idiopathic Parkinson's disease and other Parkinsonian syndromes.

Additionally, the compounds of formula Ia according to the invention may be used for the treatment and the prevention of movement disorders.

The compounds of formula Ia according to the invention may be administered
25 in conjunction with an anti-parkinsonian or any other existing therapy. For example, compounds of formula Ia may be useful as adjunct therapy in Parkinson's disease to reduce the side-effects experienced with those treatments on long term use, including but not limited to L-DOPA (motor fluctuations and dyskinesia). The compounds may also be used where in the anti-parkinsonian therapy is one of cell
30 implantation/transplantation, gene therapy, subthalamic nucleus lesions/deep brain stimulation and GPi lesion/deep brain stimulation.

The compounds of formula Ia may be used to protect against neurodegeneration and may be used in conjunction with neuroprotective agents.

The compounds of formula Ia may be used to treat neuroleptic-induced Parkinsonism and tardive dyskinesia and could be administered in conjunction with antipsychotic agents.

5 The compounds of formula Ia according to the invention may also be used for the treatment of schizophrenia, or other psychotic disorders.

The compounds of the invention may also be used in the treatment of mood disorders.

The compounds of formula Ia according to the invention may also be used in the treatment of anxiety disorders.

10 The compounds of formula Ia according to the invention may also be useful in the treatment of substance-related disorders.

The compounds of formula Ia according to the invention may also be used in the treatment of delirium, dementia, amnestic and other cognitive disorders (memory, frontal and attentional problems).

15 The compounds of formula Ia according to the invention can also be used in the treatment of sexual disorders, sleep disorders, eating disorders (anorexia/bulimia nervosa), personality disorders, factitious disorders, dissociative disorders, emesis, aggression, autism, vertigo, circadian rhythm disorders, convulsion, seizure, epilepsy, gastric motility disorders, attention deficit disorder, reward deficiency syndrome, 20 attention deficit hyperactivity disorder (ADHD), migraine, trigeminal and other neuralgias, chronic pain, neuropathic pain, cerebral ischemia, cardiac arrhythmia, myotonia, stroke, neonatal cerebral haemorrhage, amyotrophic lateral sclerosis, spasticity and degenerative diseases, bronchial asthma, asthmatic status and allergic bronchitis, asthmatic syndrome, bronchial hyperreactivity and bronchospastic 25 syndromes as well as allergic and vasomotor rhinitis and rhinoconjunctivitis.

Thus, the present invention also concerns a compound having the formula Ia or a pharmaceutically acceptable salt thereof or stereoisomeric forms thereof as defined above for use as a medicament.

30 In a further aspect, the present invention concerns also the use of a compound of formula Ia or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for the treatment of neurological, psychiatric and other disorders such as mentioned above.

In particular, the present invention concerns the use of a compound of formula Ia or a pharmaceutically acceptable salt thereof for the manufacture of a medicament 35 for the treatment of motor fluctuations and/or dyskinesia in Parkinson's patients before or during exposure to dopamine replacement therapy, idiopathic Parkinson's disease and other Parkinsonian syndromes, movement disorders, psychotic disorders,

mood disorders, anxiety disorders, substance-related disorders, delirium and dementia.

The methods of the invention comprise administration to a mammal (preferably human) suffering from above mentioned conditions or disorders, of a compound of formula Ia according to the invention in an amount sufficient to alleviate or prevent the disorder or condition.

The compound is conveniently administered in any suitable unit dosage form, including but not limited to one containing 3 to 5000 mg, preferably 5 to 500 mg of active ingredient per unit dosage form.

The term "treatment" as used herein includes symptomatic treatment, curative treatment and prophylactic treatment.

By "symptomatic" is meant the efficaciousness of the active compound in treating the current episode.

By "curative" is meant efficacy in treating the appearance of symptomatic episodes of a disorder or condition.

By "prophylactic" is meant prevention of the occurrence or recurrence of a disorder or condition. In particular we mean the prevention of any induction of the recurrent episodes and the possibility to depress the manifestation of motor fluctuation and dyskinesia.

By "motor fluctuations" is meant the development in a L-DOPA-treated subject of these four different phenomena: delayed "on" response and dose failures, end of dose wearing off, unpredictable "on-off" response and freezing episodes.

By "end-of-dose wearing off" is meant the loss of response to a dose of medication before taking the next dose.

The term "dyskinesia" is defined as the development in a subject of abnormal involuntary movements. This appears in patients with Huntington's disease, in Parkinson's disease patients exposed to chronic dopamine replacement therapy, and in Schizophrenia patients exposed to chronic treatment with neuroleptics. Dyskinesias, as a whole, are characterised by the development in a subject of abnormal involuntary movements. One way in which dyskinesias may arise is as a side effect of dopamine replacement therapy for parkinsonism or other basal ganglia-related movement disorders.

The terms "idiopathic Parkinson's disease and other Parkinsonian syndromes", include, but are not limited to genetic Parkinsonisms, multiple system atrophy, progressive supranuclear palsy, corticobasal degeneration, Fahr's disease, post-encephalitic parkinsonism, parkinsonism resulting from head injury, drug induced parkinsonisms (e.g. following treatment of schizophrenia and other psychiatric

disorders), drug intoxication (e.g. with MPTP-contaminated heroin), toxin-induced Parkinsonism (e.g. following carbon monoxide or manganese poisoning), treatment of Wilson's disease, vascular Parkinsonism, and other Parkinsonian syndromes.

The term "Parkinsonian syndrome" relates to a syndrome characterized by
5 slowness of movement (bradykinesia), rigidity and / or tremor. Parkinsonian syndromes are seen in a variety of conditions, most commonly in idiopathic Parkinsonism (i.e. Parkinson's Disease) but also following treatment of schizophrenia, exposure to toxins/drugs and head injury. It is widely appreciated that the primary pathology underlying Parkinson's disease is degeneration, in the brain, of the
10 dopaminergic projection from the substantia nigra to the striatum. This has led to the widespread use of dopamine-replacing agents (e.g. L-DOPA and dopamine agonists) as symptomatic treatments for Parkinson's disease and such treatments have been successful in increasing the quality of life of patients suffering from Parkinson's disease. However, dopamine-replacement treatments do have limitations, especially
15 following long-term treatment. Problems can include a wearing-off of the anti-parkinsonian efficacy of the treatment and the appearance of a range of side-effects which manifest as abnormal involuntary movements, such as dyskinesias.

By "movement disorder" is meant neurological motor disorders manifested by slowness or poverty of movement (bradykinesia or hypokinesia, such as that seen in
20 parkinsonian disorders) at one end of the spectrum and abnormal involuntary movement (hyperkinesias) such as tremor, dystonia, athetosis, chorea, ballismus, tics, myoclonus, restless legs syndrome, stereotypies, akathisias, and other dyskinesias at the other. Movement disorders include, but are not limited to tremors (e.g. physiological, essential, dystonic, primary writing, orthostatic, neuropathic,
25 cerebellar tremor, etc.), choreas (e.g. in Huntington's disease, Haw River syndrome, neuroacanthocytosis, McLeod syndrome, benign hereditary chorea, Sydenham's chorea, ballismus, senile chorea, etc.), tardive dyskinesia, the dystonias (e.g. childhood onset generalized primary dystonia, adult-onset primary focal and segmental dystonia, X-linked dystonia-Parkinsonism, dopa-responsive dystonia, rapid-onset dystonia, post-traumatic dystonia, tardive dystonia, paroxysmal kinesigenic
30 dyskinesia, paroxysmal nonkinesigenic dyskinesia, secondary paroxysmal dyskinesia, and other paroxysmal dyskinesias, etc.), tics including Tourette's syndrome and adult-onset tic disorders, post-infectious autoimmune neuropsychiatric disorders associated with streptococcal exposure (PANDAS), myoclonic disorders (e.g. essential,
35 posthypoxic, startle, spinal, propriospinal, toxin- and drug-induced myoclonus etc.), and other movement disorders such as hemifacial spasm, stiff person syndrome, painful legs-moving toes syndrome and restless legs syndrome.

The term "psychotic disorders" includes but is not limited to schizophrenia, schizophreniform disorder, schizoaffective disorder, delusional disorder, brief psychotic disorder, shared psychotic disorder, psychotic disorder due to a general medical condition, substance-induced psychotic disorder, postpartum psychiatric syndromes, and psychotic disorder not otherwise specified.

The term "mood disorders" includes but is not limited to depression, major depressive disorder, dysthymic disorder, depression disorder not otherwise specified, mania, bipolar I disorder, bipolar II disorder, cyclothymic disorder, bipolar disorder not otherwise specified, mood disorder due to a general medical condition, substance-induced mood disorder, mood disorder not otherwise specified,

The term "anxiety disorders" includes but is not limited to panic attack, agoraphobia, panic disorder with/without agoraphobia, agoraphobia without history of panic disorder, specific phobia, social phobia, obsessive-compulsive disorder, posttraumatic stress disorder, acute stress disorder, generalized anxiety disorder, anxiety disorder due to a general medical condition, substance-induced anxiety disorder, anxiety disorder not otherwise specified.

The term "substance-related disorders" includes more specifically substance use disorders (substance dependence, substance abuse), substance-induced disorders (substance intoxication, substance withdrawal, substance-induced mental disorders), alcohol-related disorders, amphetamine-related disorders, caffeine-related disorders, cannabis-related disorders, cocaine-related disorders, hallucinogen-related disorders, inhalant-related disorders, nicotine-related disorders, opioid related disorders, phencyclidine-related disorders, sedative-hypnotic or anxiolytic-related disorders, other (unknown) substance-related disorders.

The term "delirium" refers to a disturbance of consciousness and a change in cognition that develop over a short period of time (delirium due to a general medical condition, substance-induced delirium, delirium due to multiple etiologies, delirium not otherwise specified).

The term "dementia" refers to multiple cognitive deficits that include impairment in memory. It includes but is not limited to Alzheimer's type dementia, vascular dementia, dementia due to HIV disease, head trauma, Parkinson's disease (including dementia with Lewy bodies), Huntington's disease, Pick's disease, Creutzfeldt-Jacob disease or other medical conditions, substance-induced persisting dementia, dementia due to multiple etiologies or not otherwise specified.

The term "amnesic disorders" as used herein refers to a disturbance in memory that is either due to the direct physiological effects of a general medical condition or due to the persisting effects of a substance.

The term "epilepsy" as used herein refers to a chronic neurologic condition characterised by unprovoked, recurrent epileptic seizures. An epileptic seizure is the manifestation of an abnormal and excessive synchronised discharge of a set of cerebral neurons; its clinical manifestations are sudden and transient. The term
5 "epilepsy" as used herein can also refer to a disorder of brain function characterised by the periodic occurrence of seizures. Seizures can be "nonepileptic" when evoked in a normal brain by conditions such as high fever or exposure to toxins or "epileptic" when evoked without evident provocation.

10 The term "seizure" as used herein refers to a transient alteration of behaviour due to the disordered, synchronous, and rhythmic firing of populations of brain neurones.

The term "migraine" as used herein means a disorder characterised by recurrent attacks of headache that vary widely in intensity, frequency, and duration. The attacks are commonly unilateral and are usually associated with anorexia,
15 nausea, vomiting, phonophobia, and/or photophobia. In some cases they are preceded by, or associated with, neurological and mood disturbances. Migraine headache may last from 4 hours to about 72 hours. The International Headache Society (IHS, 1988) classifies migraine with aura (classical migraine) and migraine without aura (common migraine) as the major types of migraine. Migraine with aura consists of a headache
20 phase preceded by characteristic visual, sensory, speech, or motor symptoms. In the absence of such symptoms, the headache is called migraine without aura.

The term "bipolar disorders" as used herein refers to those disorders classified as Mood Disorders according to the Diagnostic and Statistical Manual of Mental Disorders, 4th edition (Diagnostic and Statistical Manual of Mental Disorders (DSM-IV
25 TM), American Psychiatry Association, Washington, DC, 1994). Bipolar disorders are generally characterised by spontaneously triggered repeated (i.e. at least two) episodes in which the patient's hyperexcitability, activity and mood are significantly disturbed, this disturbance consisting on some occasions of an elevation of mood and increased energy and activity (mania or hypomania), and in other occasions a lowering of mood
30 and decreased energy and activity (depression). Bipolar disorders are separated into four main categories in the DSM-IV (bipolar I disorder, bipolar II disorder, cyclothymia, and bipolar disorders not otherwise specified).

The term "manic episode", as used herein refers to a distinct period during which there is an abnormally and persistently elevated, expansive, or irritable mood
35 with signs of pressured speech and psychomotor agitation.

The term "hypomania", as used herein refers to a less extreme manic episode, with lower grade of severity.

The term "major depressive episode", as used herein refers to a period of at least 2 weeks during which there is either depressed mood or the loss of interest or pleasure in nearly all activities with signs of impaired concentration and psychomotor retardation.

5 The term "mixed episode", as used herein refers to a period of time (lasting at least 1 week) in which the criteria are met both for a manic episode and for a major depressive episode nearly every day.

10 The term "chronic pain" as used herein refers to the condition gradually being recognised as a disease process distinct from acute pain. Conventionally defined as pain that persists beyond the normal time of healing, pain can also be considered chronic at the point when the individual realises that the pain is going to be a persistent part of their lives for the foreseeable future. It is likely that a majority of chronic pain syndromes involves a neuropathic component, which is usually harder to treat than acute somatic pain.

15 The term "neuropathic pain" as used herein refers to pain due to a dysfunctional nervous system, sometimes occurring following injury to the central nervous system (central pain), but more often caused by damage to peripheral nerves (painful peripheral neuropathy). Neuropathic pain is most likely caused by neural hyperexcitation in partially damaged nerves. Several types of painful peripheral
20 neuropathy, which may share some underlying pathogenic mechanisms, have been distinguished, such as: (1) posttraumatic painful peripheral neuropathy; (2) phantom limb pain; (3) facial (trigeminal) pains; (4) postherpetic neuralgia; (5) painful diabetic neuropathy; (6) neuropathies due to cancer tumors; (7) neuropathies induced by treatment with anti-neoplastic agents; and (8) nerve damage associated with
25 demyelinating disease, such as multiple sclerosis. In neuropathic pain, painful reactions appear in response to normally neutral stimuli (allodynia) or as exaggerated reactions to painful stimuli (hyperalgesia). Spontaneous pain, not provoked by external stimuli, also occurs in neuropathic pain, and is the most difficult form of pain to measure and treat.

30 The term "tics" refers to common and often disabling neurological disorders. They are frequently associated with behaviour difficulties, including obsessive-compulsive disorder, attention deficit hyperactivity disorder and impulse control. Tics are involuntary, sudden, rapid, repetitive, nonrhythmic stereotypic movements or vocalizations. Tics are manifested in a variety of forms, with different durations and
35 degrees of complexity. Simple motor tics are brief rapid movements that often involve only one muscle group. Complex motor tics are abrupt movements that involve either a cluster of simple movements or a more coordinated sequence of movements. Simple

vocal tics include sounds such as grunting, barking, yelping, and throat clearing. Complex vocal tics include syllables, phrases, repeating other people's words and repeating one's own words.

5 The term "tremor" refers to an involuntary, rhythmical, oscillatory movement of a body part. Tremor can be phenomenologically defined as tremor at rest or associated with an action. Such an action can be postural (maintenance of a limb position), kinetic (movement-related), or intentional (at the end of a purposeful movement). Etiologically, tremor most often occurs in Parkinson's disease (Parkinsonian rest tremor) and in essential tremor (postural and kinetic tremor), which
10 consists of hereditary and age-related forms. Tremor may also occur in dystonia and in multiple sclerosis. Other tremors, which can arise from various etiologies, are cerebellar (intentional tremor) and Holmes' midbrain tremor (postural tremor). Tremor can also be an exaggerated form of normal physiological tremor. Apart from the behavioural context in which tremor occurs, tremor frequency is an important
15 criterion to distinguish between various forms of tremor. Essential tremor has the highest incidence of all tremors. As it is age-related, it can be expected to increase in aging populations. Animal models and clinical data indicate that essential tremor may be primarily based on a brainstem (inferior olivary nucleus) - cerebellar dysfunction, whereas Parkinsonian tremor probably originates from abnormal activity within the
20 basal ganglia. Excessive synchronization and/or hyperexcitation in neuronal circuits may underlie tremor activity.

The compounds of formula Ia according to the invention may advantageously be used in conjunction with one or more other therapeutic agents. In particular the compounds according to the invention may be used in conjunction with one or more
25 other therapeutic agents linked to the cholinergic transmission: e.g. agonists/antagonists to M1, M2, M3, M4, M5 receptors and to nicotinic receptors, and acetylcholinesterase modulators.

The compounds of formula Ia according to the invention may be used in conjunction with one or more other therapeutic agents linked to the
30 adrenergic/noradrenergic transmission: e.g. agonists/antagonists to $\alpha 1$, $\alpha 2$, $\beta 1$, $\beta 2$, $\beta 3$ receptors, MAO (type A and B) and COMT modulators.

The compounds of formula Ia according to the invention may be used in conjunction with one or more other therapeutic agents linked to dopaminergic transmission: e.g. agonists/antagonists to dopamine D1, D2, D3, D4, and D5
35 receptors, tyrosine-hydroxylase and DOPA-decarboxylase modulators, and vesicle monoamines transporters modulators.

The compounds of formula Ia according to the invention may be used in conjunction with one or more other therapeutic agents linked to serotonergic transmission: e.g. agonists/antagonists to 5-HT₁, 5-HT₂, 5-HT₃, 5-HT₄, 5-HT₅, 5-HT₆ and 5-HT₇ receptors.

5 The compounds of formula Ia according to the invention may be used in conjunction with one or more other therapeutic agents linked to histaminergic transmission: e.g. agonists/antagonists to H₁, H₂, H₃ and H₄ receptors.

10 The compounds of formula Ia according to the invention may be used in conjunction with one or more other therapeutic agents linked to glutamatergic transmission: e.g. agonists/antagonists to AMPA receptors (GluR1, GluR2, GluR3, GluR4); kainate receptors (GluR5, GluR6, GluR7 and KA1, KA2); NMDA receptors (subunits NR1, NR2A, NR2B, NR2C, NR2D and NR3A).

15 The compounds of formula Ia according to the invention may be used in conjunction with one or more other therapeutic agents linked to gabaergic transmission: e.g. agonists/antagonists to GABA_{A1} to GABA_{A6}, GABA_{A0}, GABA_C and GABA_B, or agents enhancing the production or reducing the degradation or the re-uptake of GABA such as vaproate, vigabatrin or tiagabine.

20 The compounds of formula Ia according to the invention may be used in conjunction with one or more other therapeutic agents such as CB₁ agonists, VR1 agonists, SV2 ligands such as levetiracetam, brivaracetam or seletracetam, amantadine, or NMDA/AMPA antagonists.

25 Activity in any of the above-mentioned indications can of course be determined by carrying out suitable clinical trials in a manner known to a person skilled in the relevant art for the particular indication and/or in the design of clinical trials in general.

For treating diseases, compounds of formula Ia or their pharmaceutically acceptable salts may be employed at an effective daily dosage and administered in the form of a pharmaceutical composition.

30 Therefore, another embodiment of the present invention concerns a pharmaceutical composition comprising an effective amount of a compound of formula Ia or pharmaceutically acceptable salts thereof in combination with a pharmaceutically acceptable diluent or carrier.

35 To prepare a pharmaceutical composition according to the invention, one or more of the compounds of formula Ia or a pharmaceutically acceptable salt thereof is intimately admixed with a pharmaceutical diluent or carrier according to conventional pharmaceutical compounding techniques known to the skilled practitioner.

Suitable diluents and carriers may take a wide variety of forms depending on the desired route of administration, e.g., oral, rectal, parenteral or intranasal.

Pharmaceutical compositions comprising compounds according to the invention can, for example, be administered orally or parenterally, i.e., intravenously,
5 intramuscularly or subcutaneously, intrathecally, by inhalation or intranasally.

Pharmaceutical compositions suitable for oral administration can be solids or liquids and can, for example, be in the form of tablets, pills, dragees, gelatin capsules, solutions, syrups, chewing-gums and the like.

To this end the active ingredient may be mixed with an inert diluent or a non-
10 toxic pharmaceutically acceptable carrier such as starch or lactose. Optionally, these pharmaceutical compositions can also contain a binder such as microcrystalline cellulose, gum tragacanth or gelatine, a disintegrant such as alginic acid, a lubricant such as magnesium stearate, a glidant such as colloidal silicon dioxide, a sweetener such as sucrose or saccharin, or colouring agents or a flavouring agent such as
15 peppermint or methyl salicylate.

The invention also contemplates compositions which can release the active substance in a controlled manner. Pharmaceutical compositions which can be used for parenteral administration are in conventional form such as aqueous or oily solutions or suspensions generally contained in ampoules, disposable syringes, glass
20 or plastics vials or infusion containers.

In addition to the active ingredient, these solutions or suspensions can optionally also contain a sterile diluent such as water for injection, a physiological saline solution, oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents, antibacterial agents such as benzyl alcohol, antioxidants such as ascorbic
25 acid or sodium bisulphite, chelating agents such as ethylene diamine-tetra-acetic acid, buffers such as acetates, citrates or phosphates and agents for adjusting the osmolarity, such as sodium chloride or dextrose.

These pharmaceutical forms are prepared using methods which are routinely used by pharmacists.

30 The amount of active ingredient in the pharmaceutical compositions can fall within a wide range of concentrations and depends on a variety of factors such as the patient's sex, age, weight and medical condition, as well as on the method of administration. Thus the quantity of compound of formula Ia in compositions for oral administration is at least 0.5 % by weight and can be up to 80 % by weight with
35 respect to the total weight of the composition.

For the preferred oral compositions, the daily dosage is in the range 5 to 5000 milligrams (mg) of compounds of formula Ia.

In compositions for parenteral administration, the quantity of compound of formula I' present is at least 0.5 % by weight and can be up to 33 % by weight with respect to the total weight of the composition. For the preferred parenteral compositions, the dosage unit is in the range 5 mg to 5000 mg of compounds of formula Ia.

The daily dose can fall within a wide range of dosage units of compound of formula Ia and is generally in the range 3 to 7000 mg, and preferably 5 to 5000 mg. However, it should be understood that the specific doses can be adapted to particular cases depending on the individual requirements, at the physician's discretion.

The following examples are provided for illustrative purposes.

Unless specified otherwise in the examples, characterization of the compounds is performed according to the following methods:

NMR spectra are recorded on a BRUKER AC 250 Fourier Transform NMR Spectrometer fitted with an Aspect 3000 computer and a 5mm $^1\text{H}/^{13}\text{C}$ dual probehead or BRUKER DRX 400 FT NMR fitted with a SG Indigo² computer and a 5 mm inverse geometry $^1\text{H}/^{13}\text{C}/^{15}\text{N}$ triple probehead. The compound is studied in DMSO- d_6 (or CDCl_3) solution at a probe temperature of 313 K or 300 K and at a concentration of 20 mg/ml. The instrument is locked on the deuterium signal of DMSO- d_6 (or CDCl_3). Chemical shifts are given in ppm downfield from TMS taken as internal standard.

HPLC analyses are performed using one of the following systems:

- an Agilent 1100 series HPLC system mounted with an INERTSIL ODS 3 C18, DP 5 μm , 250 X 4.6 mm column. The gradient runs from 100 % solvent A (acetonitrile, water, H_3PO_4 (5/95/0.001, v/v/v)) to 100 % solvent B (acetonitrile, water, H_3PO_4 (95/5/0.001, v/v/v)) in 6 min with a hold at 100 % B of 4 min. The flow rate is set at 2.5 ml/min. The chromatography is carried out at 35°C.

- a HP 1090 series HPLC system mounted with a HPLC Waters Symetry C18, 250 X 4.6 mm column. The gradient runs from 100 % solvent A (MeOH, water, H_3PO_4 (15/85/0.001M, v/v/M)) to 100 % solvent B (MeOH, water, H_3PO_4 (85/15/0.001 M, v/v/M)) in 10 min with a hold at 100 % B of 10 min. The flow rate is set at 1 ml/min. The chromatography is carried out at 40 °C.

Mass spectrometric measurements in LC/MS mode are performed as follows:

HPLC conditions

Analyses are performed using a WATERS Alliance HPLC system mounted with an INERTSIL ODS 3, DP 5 μm , 250 X 4.6 mm column.

The gradient runs from 100 % solvent A (acetonitrile, water, TFA (10/90/0.1, v/v/v)) to 100 % solvent B (acetonitrile, water, TFA (90/10/0.1, v/v/v)) in 7 min with a

hold at 100 % B of 4 min. The flow rate is set at 2.5 ml/min and a split of 1/25 is used just before API source.

MS conditions

Samples are dissolved in acetonitrile/water, 70/30, v/v at the concentration of about 250 µgr/ml. API spectra (+ or -) are performed using a FINNIGAN (San Jose, CA, USA) LCQ ion trap mass spectrometer. APCI source operates at 450 °C and the capillary heater at 160 °C. ESI source operates at 3.5 kV and the capillary heater at 210 °C.

Mass spectrometric measurements in DIP/EI mode are performed as follows: samples are vaporized by heating the probe from 50 °C to 250 °C in 5 min. EI (Electron Impact) spectra are recorded using a FINNIGAN (San Jose, CA, USA) TSQ 700 tandem quadrupole mass spectrometer. The source temperature is set at 150 °C.


Mass spectrometric measurements on a TSQ 700 tandem quadrupole mass spectrometer (Finnigan MAT, San Jose, CA, USA) in GC/MS mode are performed with a gas chromatograph model 3400 (Varian, Walnut Creek, CA, USA) fitted with a split/splitless injector and a DB-5MS fused-silica column (15 m x 0.25 mm I.D., 1 µm) from J&W Scientific (Folsom, CA, USA). Helium (purity 99.999 %) is used as carrier gas. The injector (CTC A200S autosampler) and the transfer line operate at 290 and 250 °C, respectively. Sample (1 µl) is injected in splitless mode and the oven temperature is programmed as follows: 50 °C for 5 min., increasing to 280 °C (23°C/min) and holding for 10 min. The TSQ 700 spectrometer operates in electron impact (EI) or chemical ionization (CI/CH₄) mode (mass range 33 - 800, scan time 1.00 sec). The source temperature is set at 150 °C.

Preparative chromatographic separations are performed on silicagel 60 Merck, particle size 15-40 µm, reference 1.15111.9025, using Novasep axial compression columns (80 mm i.d.), flow rates between 70 and 150 ml/min. Amount of silicagel and solvent mixtures as described in individual procedures.

The following abbreviations are used in the examples:

AcOEt	Ethyl acetate
CH ₃ CN	Acetonitrile
DMF	N,N-Dimethylformamide
NBS	N-bromosuccinimide
NCS	N-chlorosuccinimide
TFA	Trifluoroacetic acid
THF	Tetrahydrofuran

Example 1: Synthesis of 2-(7-fluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide
12.



1.1 Synthesis of 2-amino-6-fluorophenol **1**.

In a three neck flask, fitted with a magnetic stirrer, under inert atmosphere, a suspension of 2-fluoro-6-nitrophenol (5 g, 31.83 mmol), ammonium formate (6 g, 95.50 mmol) and Pd-C (10 % w/w, 0.34 g, 3.2 mmol) in MeOH (100 ml) is stirred 0.5 h at room temperature. The reaction mixture is filtered through celite and concentrated *in vacuo* to give 2-amino-6-fluorophenol **1** (4 g) which is used in the next step without any further purification.

^1H NMR ((CD_3) $_2\text{SO}$, 250 MHz): 4.66-4.98 (s (broad), 2H); 6.315 (ddd, J: 1.51; 8.10 and 10.57 Hz, 1H); 6.41 (ddd, J: 1.30, 1.30 and 8.10 Hz, 1H); 6.52 (ddd, J: 5.80, 8.10 and 8.10 Hz, 1H); 8.60-9.40 (s(broad), 1H).

1.2 Synthesis of 7-fluoro-1,3-benzoxazol-2(3H)-one **2**.

In a three neck flask, fitted with a magnetic stirrer, under inert atmosphere, 1,1-carbonyldiimidazole (15.6 g, 96 mmol) is added to a solution of 2-amino-6-fluorophenol **1** (4 g, 32 mmol) in 100 ml of anhydrous tetrahydrofuran at room temperature. The solution is refluxed for 1 h, cooled down to room temperature and concentrated *in vacuo*. The crude reaction mixture is purified by chromatography on silicagel (n-hexane/AcOEt: 1/1 (v/v)) to afford 7-fluoro-1,3-benzoxazol-2(3H)-one **2**.

Yield: 5.76 g (86 %).

^1H NMR ((CD_3) $_2\text{SO}$, 250 MHz): 6.95 (d(broad), J: 7.80 Hz, 1H); 7.03 (dd, J : 5.00 and 7.80 Hz, 1H); 7.15 (ddd, J: 5.03, 8.56 and 13.60 Hz, 1H).

Compounds listed in table 1 can be synthesized according to the same method.

Table 1:

No	IUPAC Name
3	6-methoxy-1,3-benzoxazol-2(3H)-one
4	6-fluoro-1,3-benzoxazol-2(3H)-one
5	5,6-difluoro-1,3-benzoxazol-2(3H)-one
6	7-chloro-1,3-benzoxazol-2(3H)-one
7	6,7-difluoro-1,3-benzoxazol-2(3H)-one
8	7-bromo-1,3-benzoxazol-2(3H)-one
9	7-methoxy-1,3-benzoxazol-2(3H)-one

5 1.3 Synthesis of 2-(7-fluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide **12**.


In a three neck flask, fitted with a magnetic stirrer, under inert atmosphere, NaH (in oil, 60 w/w, 1.8 g, 45.15 mmol) is carefully added by portion into a solution of 7-fluoro-1,3-benzoxazol-2(3H)-one **2** (5.76 g, 37.62 mmol) in dried DMF (80 ml) at 0 °C. After 0.25 h, 2-bromo-acetamide (6.22 g, 45.15 mmol) is added. The reaction mixture is stirred for 1 h at room temperature, quenched with water, concentrated *in vacuo* and the residue is recrystallized in EtOH/H₂O (80/20 (v/v)) to afford 2-(7-fluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide.

Yield: 3.12 g (40 %).

MS (LC-MS, MH⁺): 211.

15

Example 2: Synthesis of 2-(6-chloro-7-fluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide **19**.



20


In a three neck flask, fitted with a magnetic stirrer, under inert atmosphere, N-chlorosuccinimide (0.635 g, 4.75 mmol) is added to a suspension of 2-(7-fluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide **12** (1 g, 4.75 mmol) and sulfuric acid (90 % w/w; 20 ml) at 0 °C. The reaction is stirred 5 days then poured carefully into ice. The precipitate is collected by filtration and washed with water. The residue is

recrystallized in EtOH/water (20/80 (v/v)) to afford 0.54 g of 2-(6-chloro-7-fluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide **19**.

Yield: 0.54 g (47 %).

MS (LC-MS, MH^+): 245/247.

- 5 Example 3: Synthesis of 2-(4-chloro-6,7-difluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide **24**.



- 10 In a three neck flask, fitted with a magnetic stirrer, under inert atmosphere, N-chlorosuccinimide (0.29 g; 2.19 mmol) is added to a suspension of 2-(6,7-difluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide **18** (0.5 g, 2.19 mmol) and sulfuric acid (90 % w/w, 20 ml) at 0 °C. The reaction mixture is stirred at room temperature for two days and another portion of N-chlorosuccinimide (0.29 g, 2.19 mmol) is added. After 14 days at room temperature, the reaction mixture is poured carefully into ice. The precipitate is collected by filtration, washed with water and recrystallized EtOH/water
- 15 (20/80 (v/v)) to afford 0.42 g of a powder which is further purified by column chromatography on reverse phase to give 2-(4-chloro-6,7-difluoro-2-oxo-benzoxazol-3-yl)-acetamide **24** after recrystallisation in ethanol.

Yield: 0.163 g (30 %).

MS (LC-MS, MH^+): 261-263.

- 20 Compounds described in table 2 may be prepared according to one of the previous methods.

Table 2: Compounds of formula I.

n°	Configuration	IUPAC Name	MS(LC-MS, MH^+)
10	achiral	2-(6-methoxy-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide	223
11	achiral	2-(6-fluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide	210 (GC-MS, M^+)
12	achiral	2-(7-fluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide	211
13	achiral	2-(5,6-difluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide	229

n°	Configuration	IUPAC Name	MS(LC-MS, MH ⁺)
14	achiral	2-(6-bromo-7-fluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide	289/291
15	achiral	2-(7-chloro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide	227/229
16	achiral	2-(6-bromo-7-chloro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide	305/307/309
17	achiral	2-(6,7-dichloro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide	261/263/265
18	achiral	2-(6,7-difluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide	229
19	achiral	2-(6-chloro-7-fluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide	289/291
20	achiral	2-(7-bromo-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide	271/273
21	achiral	2-(6,7-dibromo-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide	349
22	achiral	2-(7-bromo-6-chloro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide	305/307
23	achiral	2-(7-methoxy-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide	223
24	achiral	2-(4-chloro-6,7-difluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide	263/265
25	achiral	2-(6-bromo-2-oxo-1,3-benzoxazol-3(2H)-yl)-N,N-dimethylacetamide	299/301

Example 4: pharmacological testing - hemi-parkinsonian rat model.

The invention is based upon our studies relating the use of active compounds to prolong L-DOPA activity when its action is diminishing in a rat model of Parkinson's disease.

This study was designed to investigate whether the compounds of the invention prolong L-DOPA activity using the hemi-parkinsonian rat model.

Using stereotaxic surgery in the rats, 6-hydroxydopamine (6-OHDA), a specific toxin to dopamine nerve cells, is delivered directly along the nigrostriatal pathway (medial forebrain bundle). Uptake of 6-OHDA leads to the death of dopamine nerve cells resulting in damage similar to that in PD (the abbreviation "PD" means

Parkinson's disease). In the rat, destruction of the nerve cells on one side of the brain creates an imbalance in the basal ganglia that causes the rat to spontaneously turn in a circular fashion towards the destroyed side (ipsilateral rotations). Drugs that stimulate receptor sites normally targeted by dopamine (e.g. dopamine agonist or L-DOPA) cause the rat to turn in a direction opposite to the destroyed side (contralateral rotations). The contralateral rotation response provides a behavioral index of dopamine denervation receptor supersensitivity.

The study was performed on 16 male Sprague-Dawley rats. The animals had an average weight of 270 g at surgery and 350 g during the behavioral experiments. To lesion the ascending dopaminergic nigro-striatal pathway, rats were anaesthetized with ketamine (75 mg/kg, ip) and xylazine (10 mg/kg, ip) and placed in a stereotaxic frame. 6-OHDA (10 µg/rat) was injected unilaterally into the right medial forebrain bundle.

To protect noradrenergic neurons, rats were pre-treated with imipramine (15mg/kg, ip). Two weeks after surgery, rats were challenged with 100 mg/kg i.p. L-DOPA using eight identical automated rotometers in order to select them for the test. Only rats showing at least 150 contralateral rotations within 60 min were accepted for the test. The test started one week after the selection.

The testing day, all lesioned animals were put in the testing room 15 minutes for acclimation. Eight rats were injected either with vehicle (DMSO, i.p.) or with test compound (100 µmol/kg, i.p.) 15 minutes before the L-DOPA injection and were replaced in their home cage. After the L-DOPA injection (50 mg/kg, i.p.) they were put directly into the rotometers. Contralateral rotations started to be recorded after a 10-min acclimation period in the arenas.

Analyses were conducted on the data recorded for 120 min. Data were analysed with a mixed-model analysis of variance (ANOVA) incorporating the treatment as between-group factor (treatment: 2 levels: vehicle and test compound) and the successive twelve measurements of contralateral rotations as within-subjects factor (time, 12 levels). The reliabilities of the between-mean differences within a time-sample were assessed with planned contrasts using a *F* statistic.

The major benefit of test compounds was a prolongation of L-DOPA-induced contralateral rotation during the second hour post-drug administration without rise of the side-effect (abnormal involuntary movements were not increased).

Thus the clinical benefit for test compounds may be as an adjunctive therapy to reduce motor fluctuation (i.e. "end-of-dose wearing off") and thus to increase "on-time" in parkinsonian patients exposed to dopamine-replacement therapy.


Additionally, the extension of 'on-time' and the potential for L-DOPA sparing represents a useful *de novo* therapy to delay the onset of dyskinesia.

5 The tested compounds were injected i.p. at 100 μ mol/kg. They demonstrated a significant ability to prolong L-Dopa-induced rotations in the 6-OHDA rat model during the second hour post administration (61-120 min). During this second one-hour period of the test, the following compounds induced a total number of contralateral rotations comprised between 200 and 500 turns whereas the total number of rotations observed with L-Dopa alone fell dramatically (less than 50 turns).

CLAIMS

1. A compound having formula I, or a pharmaceutically acceptable salt thereof,

5



wherein

10 R^1 is selected from hydrogen or C_{1-4} -alkyl unsubstituted or substituted by hydroxy;

R^2 is selected from hydrogen or C_{1-4} -alkyl unsubstituted or substituted by hydroxy;

R^3 is selected from hydrogen or unsubstituted C_{1-4} -alkyl;

R^{3a} is selected from hydrogen or unsubstituted C_{1-4} -alkyl;

15 R^4 is selected from hydrogen, halogen, or C_{1-4} -alkoxy unsubstituted or substituted by a phenyl group;

R^5 is selected from hydrogen, halogen, or C_{1-4} -alkoxy unsubstituted or substituted by a phenyl group;


20 R^6 is selected from hydrogen, halogen, or C_{1-4} -alkoxy unsubstituted or substituted by a phenyl group;

R^7 is selected from hydrogen, halogen, or C_{1-4} -alkoxy unsubstituted or substituted by a phenyl group;

25 with the proviso that if R^1 , R^2 , R^4 , R^5 and R^7 are hydrogen, then R^6 is not hydrogen, bromo or chloro; and if R^1 , R^2 , R^4 , R^6 and R^7 are hydrogen, then R^5 is not bromo or chloro; and if R^1 , R^2 , R^3 , R^4 and R^7 are hydrogen, R^6 is bromo and R^5 is chloro, then R^{3a} is not methyl.

2. A compound having formula I, or a pharmaceutically acceptable salt thereof,

32



wherein

R¹ is hydrogen;

R² is hydrogen;

5 R³ is selected from hydrogen or unsubstituted C₁₋₄-alkyl;

R^{3a} is selected from hydrogen or unsubstituted C₁₋₄-alkyl;

R⁴ is selected from hydrogen or halogen;

R⁵ is selected from hydrogen or halogen;

R⁶ is selected from hydrogen, halogen or unsubstituted C₁₋₄-alkyl;

10 R⁷ is selected from hydrogen, halogen or unsubstituted C₁₋₄-alkyl;

with the proviso that if R⁴, R⁵ and R⁷ are hydrogen, then R⁶ is not hydrogen, bromo or chloro; and if R⁴, R⁶ and R⁷ are hydrogen, then R⁵ is not bromo or chloro; and if R³, R⁴ and R⁷ are hydrogen, R⁶ is bromo, and R⁵ is chloro, then R^{3a} is not methyl.

15

3. A compound according to claim 1 or 2, wherein

R¹ is hydrogen;

R² is hydrogen;

R³ is selected from hydrogen or methyl;

20 R^{3a} is selected from hydrogen or methyl;

R⁴ is hydrogen or chloro;

R⁵ is hydrogen or fluoro;

R⁶ is selected from hydrogen, fluoro, chloro, bromo or methoxy;

R⁷ is selected from hydrogen, fluoro, chloro, bromo or methoxy;

25 with the proviso that if R⁴, R⁵ and R⁷ are hydrogen, then R⁶ is not hydrogen, bromo or chloro; and if R⁴, R⁶ and R⁷ are hydrogen, then R⁵ is not bromo or chloro; and if R³, R⁴ and R⁷ are hydrogen, R⁶ is bromo, and R⁵ is chloro, then R^{3a} is not methyl.

30 4. A compound according to claim 1 or 2, wherein R⁷ is halogen.

5. A compound according to claim 1 or 2, wherein R⁴ is halogen.


6. A compound selected from

- 2-(6-methoxy-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
 2-(6-fluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
 2-(7-fluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
 5 2-(5,6-difluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
 2-(6-bromo-7-fluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
 2-(7-chloro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
 2-(6-bromo-7-chloro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
 2-(6,7-dichloro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
 10 2-(6,7-difluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
 2-(6-chloro-7-fluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
 2-(7-bromo-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
 2-(6,7-dibromo-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
 2-(7-bromo-6-chloro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
 15 2-(7-methoxy-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
 2-(4-chloro-6,7-difluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
 2-(6-bromo-2-oxo-1,3-benzoxazol-3(2H)-yl)-N,N-dimethylacetamide.

7. A compound selected from

- 20 2-(6-methoxy-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
 2-(6-fluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
 2-(7-fluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
 2-(7-chloro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide;
 2-(7-bromo-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide; and
 25 2-(6,7-difluoro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide.

8. A pharmaceutical composition comprising an effective amount of a compound of formula Ia, or a pharmaceutically acceptable salt thereof,



wherein

R^1 is selected from hydrogen or C_{1-4} -alkyl unsubstituted or substituted by hydroxy;

R^2 is selected from hydrogen or C_{1-4} -alkyl unsubstituted or substituted by hydroxy;

5 R^3 is selected from hydrogen or unsubstituted C_{1-4} -alkyl;

R^{3a} is selected from hydrogen or unsubstituted C_{1-4} -alkyl;

R^4 is selected from hydrogen, halogen, or C_{1-4} -alkoxy unsubstituted or substituted by a phenyl group;


10 R^5 is selected from hydrogen, halogen, or C_{1-4} -alkoxy unsubstituted or substituted by a phenyl group;

R^6 is selected from hydrogen, halogen, or C_{1-4} -alkoxy unsubstituted or substituted by a phenyl group;

R^7 is selected from hydrogen, halogen, or C_{1-4} -alkoxy unsubstituted or substituted by a phenyl group.

15

9. A compound of formula Ia, or a pharmaceutically acceptable salt thereof,



20

wherein

R^1 is selected from hydrogen or C_{1-4} -alkyl unsubstituted or substituted by hydroxy;

R^2 is selected from hydrogen or C_{1-4} -alkyl unsubstituted or substituted by hydroxy;

25

R^3 is selected from hydrogen or unsubstituted C_{1-4} -alkyl;

R^{3a} is selected from hydrogen or unsubstituted C_{1-4} -alkyl;

R^4 is selected from hydrogen, halogen, or C_{1-4} -alkoxy unsubstituted or substituted by a phenyl group;

30

R^5 is selected from hydrogen, halogen, or C_{1-4} -alkoxy unsubstituted or substituted by a phenyl group;

R^6 is selected from hydrogen, halogen, or C_{1-4} -alkoxy unsubstituted or substituted by a phenyl group;

R⁷ is selected from hydrogen, halogen, or C₁₋₄-alkoxy unsubstituted or substituted by a phenyl group;
for use as a medicament.

5 10. A compound according to claim 9, wherein

R¹ is hydrogen;

R² is hydrogen;

R³ is selected from hydrogen or unsubstituted C₁₋₄-alkyl;

R^{3a} is selected from hydrogen or unsubstituted C₁₋₄-alkyl;

10 R⁴ is hydrogen;

R⁵ is selected from hydrogen, bromo or chloro;

R⁶ is selected from hydrogen, bromo or chloro and

R⁷ is hydrogen;

for use as a medicament.

15

11. Use of a compound according to claim 9 or 10 for the manufacture of a medicament for the treatment of the symptomatic and/or prophylactic treatment of movement disorders and/or motor fluctuations, in particular in Parkinson's disease.

20

12. Use of a compound selected from 2-(6-bromo-2-oxo-1,3-benzoxazol-3(2H)-yl)-N,N-dimethylacetamide; 2-(2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide; 2-(6-bromo-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide; 2-(6-chloro-2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide; 2-(6-chloro-2-oxo-1,3-benzoxazol-3(2H)-yl)-N,N-dimethylacetamide; and N,N-dimethyl-2-(2-oxo-1,3-benzoxazol-3(2H)-yl)acetamide for use as a medicament.

25

13. A compound selected from the group consisting of 5,6-difluoro-1,3-benzoxazol-2(3H)-one and 7-bromo-1,3-benzoxazol-2(3H)-one.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2005/005596

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07D263/58 A61K31/423 A61P25/16

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data, EMBASE, BEILSTEIN Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	KALCHEVA ET AL: "Synthesis of N-Substituted 2-Benzoxalones" CHEM. HETEROCYCL. COMPD., vol. 21, no. 9, 1985, pages 985-987, XP009052529 compound VI	8
A	WO 00/29397 A (FEENSTRA ROELOF W ; DUPHAR INT RES (NL); LONG STEPHEN K (NL); MOS JOHA) 25 May 2000 (2000-05-25) cited in the application page 1, line 18 - page 2, line 13 page 3, line 29 - page 3, line 34 ----- -/--	1-13

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- * & * document member of the same patent family

Date of the actual completion of the international search

22 August 2005

Date of mailing of the international search report

29/08/2005

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Usueli, A

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2005/005596

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 94/29272 A (ASTRA AB ; BOAR BERNARD ROBIN (GB); SHEA DENNIS MARK O (GB); TOMLINSON) 22 December 1994 (1994-12-22) cited in the application page 2, line 6 - page 3, line 16 -----	1-13
A	WO 03/076420 A (MERCK PATENT GMBH ; SCHADT OLIVER (DE); PRUECHER HELMUT (DE); LEIBROCK) 18 September 2003 (2003-09-18) cited in the application page 1, line 1 - page 4, line 13 page 5, line 7 - page 5, line 15 -----	1-13
A	EP 0 673 933 A (ADIR) 27 September 1995 (1995-09-27) cited in the application page 2, line 24 - page 3, line 11 page 6, line 9 - page 6, line 20 -----	1-13

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP2005/005596

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0029397	A	25-05-2000	AT 267183 T	15-06-2004
			AU 756470 B2	16-01-2003
			AU 1270900 A	05-06-2000
			BR 9915293 A	07-08-2001
			CA 2350137 A1	25-05-2000
			CN 1378540 A	06-11-2002
			CZ 20011657 A3	17-10-2001
			DE 69917478 D1	24-06-2004
			DE 69917478 T2	16-09-2004
			DK 1131308 T3	28-06-2004
			WO 0029397 A1	25-05-2000
			EP 1131308 A1	12-09-2001
			ES 2217833 T3	01-11-2004
			HK 1039489 A1	17-12-2004
			HU 0104922 A2	28-08-2002
			JP 3638874 B2	13-04-2005
			JP 2002530276 T	17-09-2002
			NO 20012339 A	11-05-2001
			NZ 511584 A	31-01-2003
			PL 347635 A1	22-04-2002
			RU 2225406 C2	10-03-2004
			SI 1131308 T1	31-10-2004
			SK 6492001 A3	08-10-2001
			UA 71590 C2	15-12-2004
			US 6780864 B1	24-08-2004
WO 9429272	A	22-12-1994	AU 7010894 A	03-01-1995
			CA 2164119 A1	22-12-1994
			EP 0703901 A1	03-04-1996
			FI 956074 A	18-12-1995
			JP 8511515 T	03-12-1996
			NO 955074 A	07-02-1996
			WO 9429272 A1	22-12-1994
WO 03076420	A	18-09-2003	DE 10210779 A1	09-10-2003
			AU 2003212305 A1	22-09-2003
			CA 2478945 A1	18-09-2003
			WO 03076420 A1	18-09-2003
			EP 1485363 A1	15-12-2004
			JP 2005519953 T	07-07-2005
			US 2005124627 A1	09-06-2005
EP 0673933	A	27-09-1995	FR 2717810 A1	29-09-1995
			AT 197450 T	11-11-2000
			AU 680717 B2	07-08-1997
			AU 1498895 A	28-09-1995
			CA 2145026 A1	23-09-1995
			CN 1112114 A ,C	22-11-1995
			DE 69519315 D1	14-12-2000
			DE 69519315 T2	31-05-2001
			DK 673933 T3	02-01-2001
			EP 0673933 A1	27-09-1995
			ES 2153457 T3	01-03-2001
			FI 951297 A	23-09-1995
			GR 3035175 T3	30-04-2001
			JP 2705788 B2	28-01-1998
			JP 7258233 A	09-10-1995
			NO 951081 A	25-09-1995

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2005/005596

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0673933	A	NZ 270760 A	26-03-1996
		PT 673933 T	28-02-2001
		US 5534511 A	09-07-1996
		US 5688811 A	18-11-1997
		US 5688820 A	18-11-1997
		ZA 9502333 A	15-12-1995
<hr/>			